

# The complete travel vaccine package.

It's over so quick, don't get sick

## Because you never know what's on the menu...

DISEASE	VACCINE	PATIENT AGE	DOSE	INITIAL SCHEDULE	NOTES
<b>Cholera</b>	DUKORAL® oral inactivated cholera vaccine	6+ yrs 2-6 yrs	3 mL* 3 mL†	Day 0, 7 <sup>1</sup> Day 0, 7, 14 <sup>1</sup>	>6 yrs, booster after 2 yrs 2-6 yrs, booster after 6 mths
<b>Hepatitis A</b>	AVAXIM™ inactivated hepatitis A vaccine	2+ yrs	0.5 mL	Single dose <sup>2</sup>	Booster at 6 to 12 mths
<b>Hepatitis A &amp; Typhoid Fever</b>	VIVAXIM® <i>Salmonella Typhi</i> Vi polysaccharide vaccine + inactivated hepatitis A vaccine	16+ yrs	1 mL	Single dose <sup>3</sup>	Booster of hepatitis A at 6 to 36 mths for long term protection. Revaccination against typhoid fever every 2-3 yrs
<b>Typhoid Fever</b>	TYPHIM VI™ <i>Salmonella Typhi</i> Vi polysaccharide vaccine	2+ yrs	0.5 mL	Single dose <sup>4</sup>	Revaccinate every 2-3 yrs

## ...or what may bite you,

<b>Japanese Encephalitis</b>	JE-VAX™ Japanese encephalitis vaccine	1-3 yrs 3+ yrs	0.5 mL 1.0 mL	Day 0, 7, 30 <sup>5</sup> Day 0, 7, 30	Only vaccine available against JE. Booster at 3 yr intervals <sup>12</sup>
<b>Rabies</b>	MIRV Mérieux Inactivated Rabies Vaccine	None	1.0 mL	Prophylaxis Day 0, 7, 28 <sup>6</sup>	Booster after 2 yrs <sup>12</sup>
<b>Yellow Fever</b>	STAMARIL® yellow fever vaccine	9+ mths	0.5 mL	Single dose <sup>7</sup>	Revaccinate every 10 yrs

## ...or what you can catch.

<b>Influenza Types A &amp; B</b>	VAXIGRIP® inactivated split virion influenza vaccine	3+ yrs	0.5 mL	Single dose <sup>8</sup>	VAXIGRIP® is totally thiomersal free. Yearly vaccination recommended
	VAXIGRIP® Junior	6 – 35mths	0.25mL	Single dose <sup>9</sup>	Two doses, at least 1 mth apart, are recommended for children under 9 yrs who are receiving influenza vaccine for the first time. <sup>12</sup>
<b>Poliomyelitis</b>	IPOL inactivated poliomyelitis vaccine (Vero)	2+ mths	0.5 mL	3 doses 1-2 mths apart <sup>12</sup>	Children, booster at 4 yrs of age. Adults, for travellers and those at continued risk booster every 10 years is recommended <sup>12</sup>
<b>Meningococcal Meningitis</b>	MENOMUNE® Meningococcal polysaccharide vaccine Groups A, C, Y and W-135	2+ yrs	0.5 mL	Single dose <sup>10</sup>	Revaccinate adults 3 to 5 yrs later <sup>12</sup>
<b>Diphtheria Tetanus Pertussis Poliomyelitis</b>	ADACEL®POLIO Pertussis Vaccine Acellular & Diphtheria & Tetanus Toxoids (Adsorbed) Combined with Inactivated Poliovirus Type 1, 2 and 3 (Vero cell)	4+ yrs	0.5mL	Single dose <sup>11</sup>	Use as a booster following primary vaccination

\*Mix vaccine with one sachet of buffer dissolved in 150 mL water.

†Mix vaccine with half of the buffer solution (ie. Discard 75 mL).

PBS Information: These products are not listed on the PBS except VAXIGRIP®. VAXIGRIP®: Restricted benefit – Persons at special risk of adverse consequences from infections of the lower respiratory tract.

sanofi pasteur

The vaccines business of sanofi-aventis Group